

INDIANCIVILS.COM

Online IAS Academy

Mathematics-Optional

By Venkanna Sir and Satya Sir

Partial Differential Equations 2013 - 2021

INDIANCIVILS.COM +91-9000018804

UPSC – MATHEMATICS optional – 2013 Questions

1. Form a partial differential equation by eliminating the arbitrary functions f and g from $z = yf(x) + xg(y)$. [10M]

2. Reduce the equation

$$y \frac{\partial^2 z}{\partial x^2} + (x + y) \frac{\partial^2 z}{\partial x \partial y} + x \frac{\partial^2 z}{\partial y^2} = 0$$

To its canonical form when $x \neq y$. [10M]

3. Solve $(D^2 + DD' - 6D'^2)z = x^2 \sin(x + y)$

Where D and D' denote $\frac{\partial}{\partial x}$ and $\frac{\partial}{\partial y}$. [15M]

4. Find the surface which intersects the surfaces of the system

$$z(x + y) = C(3z + 1), \text{ (C being a constant)}$$

orthogonally and which passes through the circle $x^2 + y^2 = 1, z = 1$. [15M]

5. A tightly stretched string with fixed end points $x = 0$ and $x = l$ is initially at rest in equilibrium position. If it is set vibrating by giving each point a velocity $\lambda \cdot x(l - x)$, find the displacement of the string at any distance x from one end at any time t . [20M]

UPSC – MATHEMATICS optional – 2014 Questions

1. Solve the partial differential equation $(2D^2 - 5DD' + 2D'^2)z = 24(y - x)$. [10M]

2. Reduce the equation $\frac{\partial^2 z}{\partial x^2} = x^2 \frac{\partial^2 z}{\partial y^2}$ to canonical form. [15M]

3. Find the deflection of a vibrating string (length = π , ends fixed, $\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}$) corresponding to zero initial velocity and initial deflection

$$f(x) = k(\sin x - \sin 2x) \quad [15M]$$

4. Solve $\frac{\partial^2 u}{\partial t^2} = \frac{\partial^2 u}{\partial x^2}$, $0 < x < 1, t > 0$, given that

$$(i) u(x, 0) = 0, \quad 0 \leq x \leq 1$$

$$(ii) \frac{\partial u}{\partial t}(x, 0) = x^2, \quad 0 \leq x \leq 1$$

$$(iii) u(0, t) = u(1, t) = 0, \text{ for all } t \quad [15M]$$

UPSC – MATHEMATICS optional – 2015 Questions

1. Solve the partial differential equation

$$(y^2 + z^2 - x^2)p - 2xyq + 2xz = 0$$

where $p = \frac{\partial z}{\partial x}$ and $q = \frac{\partial z}{\partial y}$. [10M]

2. Solve $(D^2 + DD' - 2D'^2)u = e^{x+y}$, where $D = \frac{\partial}{\partial x}$ and $D' = \frac{\partial}{\partial y}$. [10M]

3. Solve for the general solution $p \cos(x+y) + q \sin(x+y) = z$, where $p = \frac{\partial z}{\partial x}$ and $q = \frac{\partial z}{\partial y}$. [15M]

4. Find the solution of the initial-boundary value problem.

$$u_t - u_{xx} + u = 0, \quad 0 < x < l, t > 0$$

$$u(0, t) = u(l, t) = 0, \quad t \geq 0$$

$$u(x, 0) = x(l - x), \quad 0 < x < l$$
 [15M]

5. Reduce the second-order partial differential equation

$$x^2 \frac{\partial^2 u}{\partial x^2} - 2xy \frac{\partial^2 u}{\partial x \partial y} + y^2 \frac{\partial^2 u}{\partial y^2} + x \frac{\partial u}{\partial x} + y \frac{\partial u}{\partial y} = 0$$

Into canonical form. Hence, find its general solution. [15M]

UPSC – MATHEMATICS optional – 2016 Questions

1. Find the general equation of surfaces orthogonal to the family of spheres given by $x^2 + y^2 + z^2 = cz$. [10M]

2. Find the general integral of the partial differential equation

$$(y + zx)p - (x + yz)q = x^2 - y^2. \quad [10M]$$

3. Determine the characteristics of the equation $z = p^2 - q^2$, and find the integral surface which passes through the parabola $4z + x^2 = 0, y = 0$. [15M]

4. Solve the partial differential equation

$$\frac{\partial^3 z}{\partial x^3} - 2 \frac{\partial^3 z}{\partial x^2 \partial y} - \frac{\partial^3 z}{\partial x \partial y^2} + 2 \frac{\partial^3 z}{\partial y^3} = e^{x+y} \quad [15M]$$

5. Find the temperature $u(x, t)$ in a bar of silver of length 10 cm and constant cross-section of area 1 cm^2 . Let density $\rho = 10.6 \text{ g/cm}^3$, thermal conductivity $K = 1.04 \text{ cal/(cm sec } ^\circ\text{C)}$ and specific heat $\sigma = 0.056 \text{ cal/g } ^\circ\text{C}$. The bar is perfectly isolated laterally, with ends kept at 0°C and initial temperature $f(x) = \sin(0.1 \pi x)^\circ\text{C}$. Note that $u(x, t)$ follows the heat equation $u_t = c^2 u_{xx}$, where $c^2 = K/(\rho \sigma)$. [20M]

UPSC – MATHEMATICS optional – 2017 Questions

1. Let $f = u + iv$ be an analytic function on the unit disc $D = \{z \in \mathbb{C} : |z| < 1\}$. Show that

$$\frac{\partial^2 u}{\partial x^2} + \frac{\partial^2 u}{\partial y^2} = 0 = \frac{\partial^2 v}{\partial x^2} + \frac{\partial^2 v}{\partial y^2}$$

at all points of D.

[15M]

2. Solve $(D^2 - 2DD' + D'^2)z = e^{x+2y} + x^3 + \sin 2x$, [10M]

$$\text{Where } D \equiv \frac{\partial}{\partial x}, \quad D' = \frac{\partial}{\partial y}, \quad D^2 = \frac{\partial^2}{\partial x^2}, \quad D'^2 = \frac{\partial^2}{\partial y^2}.$$

3. Let Γ be a closed curve in xy-plane and let S denote the region bounded by the curve Γ .

Let $\frac{\partial^2 w}{\partial x^2} + \frac{\partial^2 w}{\partial y^2} = f(x, y) \forall (x, y) \in S$. If f is prescribed at each point (x, y) of S and w is prescribed on the boundary Γ of S , then prove that any solution $w = w(x, y)$, satisfying these conditions, is unique. [10M]

4. Find a complete integral of the partial differential equation

$$2(pq + yp + qx) + x^2 + y^2 = 0.$$

[15M]

5. Reduce the equation $y^2 \frac{\partial^2 z}{\partial x^2} - 2xy \frac{\partial^2 z}{\partial x \partial y} + x^2 \frac{\partial^2 z}{\partial y^2} = \frac{y^2}{x} \frac{\partial z}{\partial x} + \frac{x^2}{y} \frac{\partial z}{\partial y}$ to canonical form and hence solve it. [15M]

6. Given the one-dimensional wave equation $\frac{\partial^2 y}{\partial t^2} = c^2 \frac{\partial^2 y}{\partial x^2}; t > 0$, where $c^2 = \frac{T}{m}$, T is the constant tension in the string and m is the mass per unit length of the string.

(i) Find the appropriate solution of the above wave equation.

(ii) Find also the solution under the conditions $y(0, t) = 0$, $y(l, t) = 0$ for all t and

$$\left[\frac{\partial y}{\partial t} \right]_{t=0} = 0, y(x, 0) = a \sin \frac{\pi x}{l}, 0 < x < l. a > 0.$$

[20M]

UPSC – MATHEMATICS optional – 2018 Questions

1. Find the partial differential equation of the family of all tangent planes to the ellipsoid:
 $x^2 + 4y^2 + 4z^2 = 4$, which are not perpendicular to the xy plane [10M]

2. Solve the partial differential equation:

$$(2D^2 - 5DD' + 2D'^2)z = 5 \sin(2x + y) + 24(y - x) + e^{3x+4y}$$

Where $D \equiv \frac{\partial}{\partial x}$, $D' \equiv \frac{\partial}{\partial y}$. [15M]

3. A thin annulus occupies the region $0 < a \leq r \leq b$, $0 \leq \theta \leq 2\pi$. The faces are insulated. Along the inner edge the temperature is maintained at 0° , while along the outer edge the temperature is held at $T = K \cos \frac{\theta}{2}$, where K is a constant. Determine the temperature distribution in the annulus. [20M]

4. Find the general solution of the partial differential equation:

$(y^3x - 2x^4)p + (2y^4 - x^3y)q = 9z(x^3 - y^3)$, where $p = \frac{\partial z}{\partial x}$, $q = \frac{\partial z}{\partial y}$, and find its integral surface that passes through the curve: $x = t, y = t^2, z = 1$. [15M]

UPSC – MATHEMATICS optional – 2019 Questions

1. Form a partial differential equation of the family of surfaces given by the following expression: $\psi(x^2 + y^2 + 2z^2, y^2 - 2zx) = 0$. [10M]

2. Solve the first order quasilinear partial differential equation by the method of characteristics: [15M]

$$x \frac{\partial u}{\partial x} + (u - x - y) \frac{\partial u}{\partial y} = x + 2y \text{ in } x > 0, -\infty < y < \infty \text{ with } u = 1 + y \text{ on } x = 1.$$

3. Reduce the following second order partial differential equation to canonical form and find the general solution:

$$\frac{\partial^2 u}{\partial x^2} - 2x \frac{\partial^2 u}{\partial x \partial y} + x^2 \frac{\partial^2 u}{\partial y^2} = \frac{\partial y}{\partial x} + 12x. [20M]$$

UPSC – MATHEMATICS optional – 2020 Questions

1. Form a partial differential equation by eliminating the arbitrary functions $f(x)$ and $g(y)$ from $z = y f(x) + x g(y)$ and specify its nature (elliptic, hyperbolic or parabolic in the region $x > 0, y > 0$). [10M]

2. Solve the partial differential equation:

$$(D^3 - 2D^2D' - DD'^2 + 2D'^3)z = e^{2x+y} + \sin(x - 2y);$$
$$D \equiv \frac{\partial}{\partial x}, D' \equiv \frac{\partial}{\partial y} [10M]$$

3. Find the integral surface of the partial differential equation:

$$(x - y)y^2 \frac{\partial z}{\partial x} + (y - x)x^2 \frac{\partial z}{\partial y} = (x^2 + y^2)z \text{ that contains the curve: } xz = a^3, y = 0, \text{ on it.}$$

[15M]

4. Find the solution of the partial differential equation: $z = \frac{1}{2}(p^2 + q^2) + (p - x)(q - y)$;
 $p \equiv \frac{\partial z}{\partial x}$, $q \equiv \frac{\partial z}{\partial y}$ which passes through the x -axis.

[15M]

UPSC – MATHEMATICS optional – 2021 Questions

1. Obtain the partial differential equation by eliminating arbitrary function f from the equation $f(x + y + z, x^2 + y^2 + z^2) = 0$

[10M]

2. Solve the wave equation $a^2 \frac{\partial^2 u}{\partial x^2} = \frac{\partial^2 u}{\partial t^2}$, $0 < x < L$, $t > 0$

Subject to the conditions

$$u(0, t) = 0, \quad u(L, t) = 0 \quad u(x, 0) = \frac{1}{4}x(L - x), \quad \left. \frac{\partial u}{\partial t} \right|_{t=0} = 0$$

[10M]

3. Find the general solution of the partial differential equation

$$(D^2 - D'^2 - 3D + 3D')z = xy + e^{x+2y} \text{ Where } D \equiv \frac{\partial}{\partial x} \text{ and } D' \equiv \frac{\partial}{\partial y}. \quad [15M]$$

4. Find a complete integral of the partial differential equation $p = (z + qy)^2$ by using Charpit's method.

[15M]